

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/66, 31/675	A1	(11) International Publication Number: WO 95/30421 (43) International Publication Date: 16 November 1995 (16.11.95)
(21) International Application Number: PCT/IB95/00288 (22) International Filing Date: 24 April 1995 (24.04.95) (30) Priority Data: 9408775.6 4 May 1994 (04.05.94) GB (71) Applicant (for all designated States except US): CIBA-GEIGY AG [CH/CH]; Klybeckstrasse 141, CH-4002 Basle (CH). (72) Inventors; and (75) Inventors/Applicants (for US only): GOODSHIP, Allen, E. [GB/GB]; "Dormers", Mead Road, Stoke Gifford, Bristol BS12 6PS (GB). KENWRIGHT, John [GB/GB]; The Farm House, Cutts End, Cumnor, Oxford OX2 9QH (GB). GREEN, Jonathan [GB/CH]; Waldstrasse 12, CH-4144 Arlesheim (CH). (74) Common Representative: CIBA-GEIGY AG; Patentabteilung, Klybeckstrasse 141, CH-4002 Basle (CH).		(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, IS, JP, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). Published <i>With international search report.</i>
(54) Title: USE OF CERTAIN METHANE BISPHOSPHONIC ACID DERIVATIVES TO PREVENT PROSTHESIS LOOSENING AND PROSTHESIS MIGRATION (57) Abstract The invention relates to the use of certain methanebisphosphonic acid derivatives (for the manufacture of a pharmaceutical composition) for the prevention and treatment of prosthesis loosening and prosthesis migration in mammals including man.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Use of certain methanebisphosphonic acid derivatives to prevent prosthesis loosening and prosthesis migration

Arthroplasty, especially of the hip joint, is now a common procedure for the treatment of patients with osteoarthritis, rheumatoid arthritis and osteoporotic fracture. Approximately 800 000 hip replacements are performed globally each year. This number is rising steadily, younger patients are being treated and patients are living much longer and they all expect to lead a mobile, independent life. Despite the indisputable success of this technique, often complications arise and may eventually require costly surgical revision (3 or 4 revisions being not uncommon for a single patient).

The most common complications are aseptic prosthesis loosening [see for example H. Malchau et al., *Acta Orthop. Scand.* 64 (1993) 497-506; R.D. Mulroy Jr. and W.H. Harris, *J. Bone Joint Surg. (Br)* 72-B (1990) 757-760; M. Winter et al., *Clin. Orthop. Rel. Research* 282 (1992) 73-80] and migration of the prosthesis through bone without overt loosening [see for example B.B. Wroblewski and P.D. Siney, *Clin. Orthop. Rel. Research* 285 (1992) 45-47; M. Winter et al., *loc. cit.*]. These complications result in considerable morbidity, such as pain and reduced mobility. They occur with both cemented and uncemented prostheses, regardless of design, material or coating.

Previous approaches to prevent these complications have concentrated on alterations to the design of the prosthesis and the selection of different materials for its construction [see for example H. Malchau et al., *loc. cit.*; R.D. Mulroy Jr. and W.H. Harris, *loc. cit.*; M. Winter et al., *loc. cit.*; B.B. Wroblewski and P.D. Siney, *loc. cit.*; J. Wilson-MacDonald et al., *J. Bone Joint Surg. (Br)* 72-B (1990) 423-430]. Apart from some preliminary research with non-steroidal anti-inflammatory drugs [A. Ohlin and U.H. Lerner, *Bone and Mineral* 20 (1993) 67-78; J.H. Herman et al., *J. Rheumatol.* 21 (1994) 338-343], a pharmacological approach to prevent or treat the complications of arthroplasty, especially hip arthroplasty, has not been successful.

Methanebisphosphonic acid derivatives, in particular bisphosphonate compounds (= bisphosphonates), are used clinically to inhibit excessive bone resorption in a variety of diseases such as tumour-induced osteolysis, Paget's disease and osteoporosis [H. Fleisch, *Handbook of Exptl. Pharmacol.* 83 (1988) 441-465]. Radiolabelled bisphosphonates are also used diagnostically to identify sites of high bone turnover [B.D. Collier et al., *J. Nucl. Med.* 34 (1993) 2241-2246; I. Fogelman et al., *J. Nucl. Med.* 34 (1993) 2247-2252].

In the orthopaedic field in general, there is a strong prejudice against using bisphosphonates. Firstly, attempts to use the commercially available bisphosphonate etidronate in the prevention of heterotopic ossification - which is another complication after arthroplasty but is very different from prosthesis loosening and prosthesis migration - proved either ineffective, or even detrimental due to a rebound effect [see for example D.E. Garland, Clin. Orthop. Rel. Research 263 (1991) 13-29; B.J. Thomas and H.C. Amstutz, The Hip 1987, 59-69; B.J. Thomas, Orthop. Clin. North America 23 (1992) 347-358].

Furthermore, there is increasing evidence that some bisphosphonates may inhibit bone formation and mineralization. Kanis, for example, teaches in Lancet 1984, 27-33 that bisphosphonates, as inhibitors of bone resorption, may delay the repair of microfractures by reducing the rate of remodelling. As another example, Reid et al. in Lancet 1988, 143-146 found that bisphosphonate treatment, in this particular case done with disodium pamidronate, caused a reduction in bone formation and a very low rate of bone turnover which raised the possibility of impaired microfracture repair. Adamson et al. [Lancet 342 (1993) 1459-1460] reports that pamidronate may inhibit bone mineralization in Paget's disease.

Surprisingly, it has now been found that certain methanebisphosphonic acid derivatives are useful for the cost-effective prevention or treatment of these complications of joint replacement, especially hip replacement, in mammals including man.

This can be demonstrated by appropriate in vitro and in vivo tests. In particular, in vivo experiments show that the bisphosphonates according to the invention are effective in preventing prosthesis loosening in the treated animals, e.g. sheep. This is demonstrated, for example, by the following test methods: 12 adult female sheep, approx. 50 kg body weight, are randomly divided into 2 groups of 6 animals and all undergo a unilateral total hip replacement operation with implantation of a titanium prosthesis. Immediately after surgery one group of animals receives an i.v. infusion of bisphosphonate and the other group receives saline. This treatment may be subsequently repeated, for example at 4 and 8 weeks. At regular monthly intervals the hip and femoral region of each animal is subjected to both standard and quantitative digital radiography to assess changes in bone density around the prosthesis as an early indicator of prosthesis loosening. In addition the animals are walked over a Kistler force plate to measure vertical ground reaction force

from both hind limbs to determine limb functionality and, indirectly, pain. At 12 months after the operation, and 2 weeks prior to sacrifice, the animals receive a double injection, 10 days apart, of the fluorescent bone label oxytetracycline to facilitate histomorphometric analysis. At the end of the labelling period the animals are killed and the femurs with the implants are removed for post-mortem analysis. Precision sections, cut and polished from the femurs, are subjected to microradiography to determine bone loss around the implant. Similar sections are also analyzed by histomorphometry to assess bone resorption and formation at the tissue level. Finally sections of bone containing the implant are subjected to a mechanical "push-out" test on a materials testing machine to measure the strength of bonding between the metal implant and the surrounding bone.

A comparison of the test animals treated with a bisphosphonate according to the invention with the controls clearly shows that bisphosphonate treatment is beneficial to prevent and treat prosthesis loosening.

The invention therefore relates to the use of a methanebisphosphonic acid derivative selected from

3-amino-1-hydroxypropane-1,1-diphosphonic acid (pamidronic acid), e.g. pamidronate; 3-(N,N-dimethylamino)-1-hydroxypropane-1,1-diphosphonic acid, e.g. dimethyl-APD; 4-amino-1-hydroxybutane-1,1-diphosphonic acid (alendronic acid), e.g. alendronate; 6-amino-1-hydroxyhexane-1,1-diphosphonic acid, e.g. amino-hexyl-BP; 3-(N-methyl-N-n-pentylamino)-1-hydroxypropane-1,1-diphosphonic acid, e.g. methyl-pentyl-APD (= BM 21.0955); 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid; 1-hydroxy-2-(3-pyridyl)ethane-1,1-diphosphonic acid (risedronic acid), e.g. risedronate, including N-methyl pyridinium salts thereof, for example N-methyl pyridinium iodides such as NE-10244 or NE-10446; 1-(4-chlorophenylthio)methane-1,1-diphosphonic acid (tiludronic acid), e.g. tiludronate; 3-[N-(2-phenylthioethyl)-N-methylamino]-1-hydroxypropane-1,1-diphosphonic acid; 1-hydroxy-3-(pyrrolidin-1-yl)propane-1,1-diphosphonic acid, e.g. EB 1053 (Leo); 1-(N-phenylaminothiocarbonyl)methane-1,1-diphosphonic acid, e.g. FR 78844 (Fujisawa); 5-benzoyl-3,4-dihydro-2H-pyrazole-3,3-diphosphonic acid tetraethyl ester, e.g. U-81581 (Upjohn); 1-hydroxy-2-(imidazo[1,2-a]pyridin-3-yl)ethane-1,1-diphosphonic acid, e.g. YM 529; and 1,1-dichloromethane-1,1-diphosphonic acid (clodronic acid), e.g. clodronate; or a pharmaceutically acceptable salt thereof, or any hydrate thereof,

(for the manufacture of a pharmaceutical composition) for the prevention and treatment of prosthesis loosening and prosthesis migration in mammals including man.

As already mentioned above, prosthesis loosening and prosthesis migration often occurs as a complication after arthroplasty. Arthroplasty means a partial, e.g. hemi, or total arthroplasty of any joint, especially hip and knee, and in particular hip.

Pharmaceutically acceptable salts are preferably salts with bases, conveniently metal salts derived from groups Ia, Ib, IIa and IIb of the Periodic Table of the Elements, including alkali metal salts, e.g. potassium and especially sodium salts, or alkaline earth metal salts, preferably calcium or magnesium salts, and also ammonium salts with ammonia or organic amines.

Especially preferred pharmaceutically acceptable salts are those where one, two, three or four, in particular one or two, of the acidic hydrogens of the bisphosphonic acid are replaced by a pharmaceutically acceptable cation, in particular sodium, potassium or ammonium, in first instance sodium.

A very preferred group of pharmaceutically acceptable salts is characterized by having one acidic hydrogen and one pharmaceutically acceptable cation, especially sodium, in each of the phosphonic acid groups.

All the methanebisphosphonic acid derivatives mentioned above are well known from the literature. This includes their manufacture (see e.g. EP-A-513 760, pp. 13-48). For example, 3-amino-1-hydroxypropane-1,1-diphosphonic acid is prepared as described e.g. in US patent 3,962,432 as well as in US patents 4,639,338 and 4,711,880, and 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid is prepared as described e.g. in US patent 4,939,130.

A particular embodiment of the invention is represented by the use of a methane-bisphosphonic acid derivative which is selected from 3-amino-1-hydroxypropane-1,1-diphosphonic acid, 3-(N,N-dimethylamino)-1-hydroxypropane-1,1-diphosphonic acid, 4-amino-1-hydroxybutane-1,1-diphosphonic acid, 6-amino-1-hydroxyhexane-1,1-diphosphonic acid, 3-(N-methyl-N-n-pentylamino)-1-hydroxypropane-1,1-diphosphonic acid, 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid; 1-hydroxy-2-(3-pyridyl)ethane-1,1-diphosphonic acid, and N-methyl pyridinium salts thereof; 1-(4-chloro-

phenylthio)methane-1,1-diphosphonic acid, 3-[N-(2-phenylthioethyl)-N-methylamino]-1-hydroxypropane-1,1-diphosphonic acid; 1-hydroxy-3-(pyrrolidin-1-yl)propane-1,1-diphosphonic acid, 1-(N-phenylaminothiocarbonyl)methane-1,1-diphosphonic acid, 5-benzoyl-3,4-dihydro-2H-pyrazole-3,3-diphosphonic acid tetraethyl ester, 1-hydroxy-2-(imidazo[1,2-a]pyridin-3-yl)ethane-1,1-diphosphonic acid, a pharmaceutically acceptable salt thereof, and any hydrate thereof.

A preferred embodiment of the invention is represented by the use of a methane-bisphosphonic acid derivative which is selected from 3-amino-1-hydroxypropane-1,1-diphosphonic acid, 3-(N,N-dimethylamino)-1-hydroxypropane-1,1-diphosphonic acid, 4-amino-1-hydroxybutane-1,1-diphosphonic acid, 6-amino-1-hydroxyhexane-1,1-diphosphonic acid, 3-(N-methyl-N-n-pentylamino)-1-hydroxypropane-1,1-diphosphonic acid, 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid; 1-hydroxy-2-(3-pyridyl)ethane-1,1-diphosphonic acid, 3-[N-(2-phenylthioethyl)-N-methylamino]-1-hydroxypropane-1,1-diphosphonic acid; 1-hydroxy-3-(pyrrolidin-1-yl)propane-1,1-diphosphonic acid, 1-hydroxy-2-(imidazo[1,2-a]pyridin-3-yl)ethane-1,1-diphosphonic acid, a pharmaceutically acceptable salt thereof, and any hydrate thereof.

A very preferred embodiment of the invention is represented by the use of a methane-bisphosphonic acid derivative which is selected from pamidronic acid, alendronic acid, 3-(N-methyl-N-n-pentylamino)-1-hydroxypropane-1,1-diphosphonic acid, 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid, risedronic acid and tiludronic acid, a pharmaceutically acceptable salt thereof, and any hydrate thereof.

An especially preferred embodiment of the invention is represented by the use of a methanebisphosphonic acid derivative which is selected from 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid and 3-amino-1-hydroxypropane-1,1-diphosphonic acid, a pharmaceutically acceptable salt thereof, and any hydrate thereof.

Further the invention relates to the use of 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid or a pharmaceutically acceptable salt thereof or any hydrate thereof.

Further the invention relates to the use of 3-amino-1-hydroxypropane-1,1-diphosphonic acid or a pharmaceutically acceptable salt thereof or any hydrate thereof.

The pharmaceutically useful methanebisphosphonic acid derivatives may be used in the

form of a possible isomer or of a mixture of isomers, typically as optical isomers such as enantiomers or diastereoisomers or geometric isomers, typically cis-trans isomers. The optical isomers are obtained in the form of the pure antipodes and/or as racemates.

The pharmaceutically useful methanebisphosphonic acid derivatives can also be used in the form of their hydrates or include other solvents used for their crystallisation.

The methanebisphosphonic acid derivative is preferably used in the form of a pharmaceutical preparation that contains a therapeutically effective amount of the active ingredient optionally together with or in admixture with inorganic or organic, solid or liquid, pharmaceutically acceptable carriers which are suitable for administration.

The pharmaceutical compositions may be, for example, compositions for enteral, such as oral, rectal, aerosol inhalation or nasal administration, compositions for parenteral, such as intravenous or subcutaneous administration, or compositions for transdermal administration (e.g. passive or iontophoretic). Furthermore, direct local administration at the time of operation comes into consideration. What is meant by the latter type of administration is the coating of the exposed bone surface to which the implant is attached, with a bisphosphonate according to the invention, for example by washing the bone surface with a bisphosphonate solution. What is not meant by this type of administration is the coating of the prosthesis with bisphosphonates.

Preferably, the pharmaceutical compositions are adapted to oral or parenteral (especially intravenous or transdermal) administration before, during or after prosthesis implantation, or to local administration during prosthesis implantation. Intravenous and oral, first and foremost intravenous, administration is considered to be of particular importance.

The particular mode of administration and the dosage will be selected by the attending physician taking into account the particulars of the patient, especially age, weight, life style, activity level, hormonal status (e.g. post-menopausal), bone mineral density and type of prosthesis to be implanted.

The dosage of the active ingredient may depend on various factors, such as effectiveness and duration of action of the active ingredient, mode of administration, warm-blooded species, and/or sex, age, weight and individual condition of the warm-blooded animal.

Normally the dosage is such that a single dose of from 0.002 - 3.40 mg/kg, especially 0.01 - 2.40 mg/kg, is administered to a warm-blooded animal weighing approximately 75 kg. If desired, this dose may also be taken in several, optionally equal, partial doses.

"mg/kg" means mg drug per kg body weight of the mammal - including man - to be treated.

The dose mentioned above - either administered as a single dose (which is preferred) or in several partial doses - may be repeated, for example once daily, once weekly, once every month, once every three months, ~~once every six months~~ or once a year. In other words, the pharmaceutical compositions may be administered in regimens ranging from continuous daily therapy to intermittent cyclical therapy.

Preferably, the methanebisphosphonic acid derivatives are administered in doses which are in the same order of magnitude as those used in the treatment of the diseases classically treated with methanebisphosphonic acid derivatives, such as Paget's disease, tumour-induced hypercalcaemia or osteoporosis. In other words, preferably the methanebisphosphonic acid derivatives are administered in doses which would likewise be therapeutically effective in the treatment of Paget's disease, tumour-induced hypercalcaemia or osteoporosis, i.e. preferably they are administered in doses which would likewise effectively inhibit bone resorption.

Formulations in single dose unit form contain preferably from about 1 % to about 90 %, and formulations not in single dose unit form contain preferably from about 0.1 % to about 20 %, of the active ingredient. Single dose unit forms such as capsules, tablets or dragées contain e.g. from about 1 mg to about 500 mg of the active ingredient.

The invention further relates to the use of a composition for the manufacture of a medicament, e.g. in single dose unit form, for the prevention and treatment of prosthesis loosening and prosthesis migration in mammals including man, wherein the composition contains 0.002 - 3.40 mg/kg, especially 0.01 - 2.40 mg/kg, of a methanebisphosphonic acid derivative as defined above per dosage form.

Moreover, the invention relates to the use of a methanebisphosphonic acid derivative as defined above at a dose of 0.002 - 3.40 mg/kg, especially 0.01 - 2.40 mg/kg, per dosage form for the manufacture of a medicament for the prevention and treatment of prosthesis

loosening and prosthesis migration in mammals including man.

Pharmaceutical preparations for enteral and parenteral administration are, for example, those in dosage unit forms, such as dragées, tablets or capsules and also ampoules. They are prepared in a manner known per se, for example by means of conventional mixing, granulating, confectioning, dissolving or lyophilising processes. For example, pharmaceutical preparations for oral administration can be obtained by combining the active ingredient with solid carriers, where appropriate granulating a resulting mixture, and processing the mixture or granulate, if desired or necessary after the addition of suitable adjuncts, into tablets or dragée cores.

Suitable carriers are especially fillers, such as sugars, for example lactose, saccharose, mannitol or sorbitol, cellulose preparations and/or calcium phosphates, for example tricalcium phosphate or calcium hydrogen phosphate, and also binders, such as starch pastes, using, for example, corn, wheat, rice or potato starch, gelatin, tragacanth, methylcellulose and/or polyvinylpyrrolidone and, if desired, disintegrators, such as the above-mentioned starches, also carboxymethyl starch, crosslinked polyvinylpyrrolidone, agar or alginic acid or a salt thereof, such as sodium alginate. Adjuncts are especially flow-regulating agents and lubricants, for example silicic acid, talc, stearic acid or salts thereof, such as magnesium or calcium stearate, and/or polyethylene glycol. Drage cores are provided with suitable coatings that may be resistant to gastric juices, there being used, inter alia, concentrated sugar solutions that optionally contain gum arabic, talc, polyvinylpyrrolidone, polyethylene glycol and/or titanium dioxide, or lacquer solutions in suitable organic solvents or solvent mixtures or, to produce coatings that are resistant to gastric juices, solutions of suitable cellulose preparations, such as acetylcellulose phthalate or hydroxypropylmethylcellulose phthalate. Colouring substances or pigments may be added to the tablets or drage coatings, for example for the purpose of identification or to indicate different doses of active ingredient.

Other orally administrable pharmaceutical preparations are dry-filled capsules made of gelatin, and also soft, sealed capsules made of gelatin and a plasticiser, such as glycerol or sorbitol. The dry-filled capsules may contain the active ingredient in the form of a granulate, for example in admixture with fillers, such as lactose, binders, such as starches, and/or glidants, such as talc or magnesium stearate, and, where appropriate, stabilisers. In soft capsules the active ingredient is preferably dissolved or suspended in suitable liquids, such as fatty oils, paraffin oil or liquid polyethylene glycols, it being possible also for

stabilisers to be added.

Parenteral formulations are especially injectable fluids that are effective in various manners, such as intravenously, intramuscularly, intraperitoneally, intranasally, intradermally or subcutaneously. Such fluids are preferably isotonic aqueous solutions or suspensions which can be prepared before use, for example from lyophilised preparations which contain the active ingredient alone or together with a pharmaceutically acceptable carrier. The pharmaceutical preparations may be sterilised and/or contain adjuncts, for example preservatives, stabilisers, wetting agents and/or emulsifiers, solubilisers, salts for regulating the osmotic pressure and/or buffers.

Suitable formulations for transdermal application include an effective amount of the active ingredient with carrier. Advantageous carriers include absorbable pharmacologically acceptable solvents to assist passage through the skin of the host. Characteristically, transdermal devices are in the form of a bandage comprising a backing member, a reservoir containing the compound optionally with carriers, optionally a rate controlling barrier to deliver the active ingredient of the skin of the host at a controlled and predetermined rate over a prolonged period of time, and means to secure the device to the skin.

The following Examples illustrate the invention described hereinbefore. The term "active ingredient" is to be understood as being any one of the methanebisphosphonic acid derivatives mentioned above as being useful according to the present invention.

Example 1: Capsules containing coated pellets of, for example, disodium pamidronate pentahydrate, as active ingredient:

Core pellet:

active ingredient (ground)	197.3 mg
Microcrystalline cellulose	52.7 mg
(Avicel® PH 105)	—
	250.0 mg

+ Inner coating:

Cellulose HP-M 603	10.0 mg
Polyethylene glycol	2.0 mg

- 10 -

Talc	8.0 mg
	<hr/>
	270.0 mg
+ Gastric juice-resistant outer coating:	
Eudragit® L 30 D (solid)	90.0 mg
Triethyl citrate	21.0 mg
Antifoam® AF	2.0 mg
Water	
Talc	7.0 mg
	<hr/>
	390.0 mg

A mixture of disodium pamidronate with Avicel® PH 105 is moistened with water and kneaded, extruded and formed into spheres. The dried pellets are then successively coated in the fluidized bed with an inner coating, consisting of cellulose HP-M 603, polyethylene glycol (PEG) 8000 and talc, and the aqueous gastric juice-resistant coat, consisting of Eudragit® L 30 D, triethyl citrate and Antifoam® AF. The coated pellets are powdered with talc and filled into capsules (capsule size 0) by means of a commercial capsule filling machine, for example Höfliger and Karg.

Example 2: Monolith adhesive transdermal system, containing as active ingredient, for example, 1-hydroxy-2-(imidazol-1-yl)-ethane-1,1-diphosphonic acid:

Composition:

polyisobutylene (PIB) 300 (Oppanol B1, BASF)	5.0 g
PIB 35000 (Oppanol B10, BASF)	3.0 g
PIB 1200000 (Oppanol B100, BASF)	9.0 g
hydrogenated hydrocarbon resin (Escorez 5320, Exxon)	43.0 g
1-dodecylazacycloheptan-2-one (Azone, Nelson Res., Irvine/CA)	20.0 g

- 11 -

active ingredient	20.0 g
Total	100.0 g

Preparation:

The above components are together dissolved in 150 g of special boiling point petroleum fraction 100-125 by rolling on a roller gear bed. The solution is applied to a polyester film (Hostaphan, Kalle) by means of a spreading device using a 300 μm doctor blade, giving a coating of about 75 g/m^2 . After drying (15 minutes at 60°C), a silicone-treated polyester film (thickness 75 μm , Laufenberg) is applied as the peel-off film. The finished systems are punched out in sizes in the wanted form of from 5 to 30 cm^2 using a punching tool. The complete systems are sealed individually in sachets of aluminised paper.

Example 3: Vial containing 1.0 mg dry, lyophilized 1-hydroxy-2-(imidazol-1-yl)-ethane-1,1-diphosphonic acid (mixed sodium salts thereof). After dilution with 1 ml of water, a solution (concentration 1 mg/ml) for i.v. infusion is obtained.

Composition:

active ingredient (free diphosphonic acid)	1.0 mg
mannitol	46.0 mg
Trisodium citrate x 2 H ₂ O	ca. 3.0 mg
water	1 ml

water for injection	1 ml .
---------------------	--------

In 1 ml of water, the active ingredient is titrated with trisodium citrate x 2 H₂O to pH 6.0. Then, the mannitol is added and the solution is lyophilized and the lyophilisate filled into a vial.

Example 4: Ampoule containing disodium pamidronate pentahydrate dissolved in water. The solution (concentration 3-mg/ml) is for i.v. infusion after dilution.

Composition:

active ingredient	19.73 mg
-------------------	----------

(\approx 15.0 mg of anhydrous active ingredient)

mannitol	250 mg
water for injection	5 ml .

Example 5: Tablets each containing 50 mg of 3-[N-(2-phenylthioethyl)-N-methylamino]-1-hydroxypropane-1,1-diphosphonic acid can be prepared as follows:

Composition (10,000 tablets)

Active ingredient	500.0 g
Lactose	500.0 g
Potato starch	325.0 g
Gelatin	8.0 g
Talc	60.0 g
Magnesium stearate	10.0 g
Silicon dioxide (finely divided)	20.0 g
Ethanol	q.s.

The active ingredient is mixed with the lactose and 292 g of potato starch, and the mixture is moistened with an ethanolic solution of the gelatin and granulated through a sieve. After the granules have dried, the remainder of the potato starch, the magnesium stearate and the silicon dioxide are admixed and the mixture compressed to give tablets each weighing 145.0 mg and containing 50.0 mg of active ingredient, which can, if desired, be provided with breaking grooves to enable the dosage to be more finely adjusted.

Claims:

1. Use of a methanebisphosphonic acid derivative selected from

3-amino-1-hydroxypropane-1,1-diphosphonic acid, 3-(N,N-dimethylamino)-1-hydroxypropane-1,1-diphosphonic acid, 4-amino-1-hydroxybutane-1,1-diphosphonic acid, 6-amino-1-hydroxyhexane-1,1-diphosphonic acid, 3-(N-methyl-N-n-pentylamino)-1-hydroxypropane-1,1-diphosphonic acid, 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid; 1-hydroxy-2-(3-pyridyl)ethane-1,1-diphosphonic acid, and N-methyl pyridinium salts thereof; 1-(4-chlorophenylthio)methane-1,1-diphosphonic acid, 3-[N-(2-phenylthioethyl)-N-methylamino]-1-hydroxypropane-1,1-diphosphonic acid; 1-hydroxy-3-(pyrrolidin-1-yl)propane-1,1-diphosphonic acid, 1-(N-phenylaminothiocarbonyl)methane-1,1-diphosphonic acid, 5-benzoyl-3,4-dihydro-2H-pyrazole-3,3-diphosphonic acid tetraethyl ester, 1-hydroxy-2-(imidazo[1,2-a]pyridin-3-yl)ethane-1,1-diphosphonic acid and 1,1-dichloromethane-1,1-diphosphonic acid; or a pharmaceutically acceptable salt thereof, or any hydrate thereof,

for the manufacture of a pharmaceutical composition for the prevention and treatment of prosthesis loosening and prosthesis migration in mammals including man.

2. Use according to claim 1 of a methanebisphosphonic acid derivative which is selected from 3-amino-1-hydroxypropane-1,1-diphosphonic acid, 3-(N,N-dimethylamino)-1-hydroxypropane-1,1-diphosphonic acid, 4-amino-1-hydroxybutane-1,1-diphosphonic acid, 6-amino-1-hydroxyhexane-1,1-diphosphonic acid, 3-(N-methyl-N-n-pentylamino)-1-hydroxypropane-1,1-diphosphonic acid, 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid; 1-hydroxy-2-(3-pyridyl)ethane-1,1-diphosphonic acid, and N-methyl pyridinium salts thereof; 1-(4-chlorophenylthio)methane-1,1-diphosphonic acid, 3-[N-(2-phenylthioethyl)-N-methylamino]-1-hydroxypropane-1,1-diphosphonic acid; 1-hydroxy-3-(pyrrolidin-1-yl)propane-1,1-diphosphonic acid, 1-(N-phenylaminothiocarbonyl)methane-1,1-diphosphonic acid, 5-benzoyl-3,4-dihydro-2H-pyrazole-3,3-diphosphonic acid tetraethyl ester, 1-hydroxy-2-(imidazo[1,2-a]pyridin-3-yl)ethane-1,1-diphosphonic acid, a pharmaceutically acceptable salt thereof, and any hydrate thereof.

3. Use according to claim 1 of a methanebisphosphonic acid derivative which is selected

from 3-amino-1-hydroxypropane-1,1-diphosphonic acid, 3-(N,N-dimethylamino)-1-hydroxypropane-1,1-diphosphonic acid, 4-amino-1-hydroxybutane-1,1-diphosphonic acid, 6-amino-1-hydroxyhexane-1,1-diphosphonic acid, 3-(N-methyl-N-n-pentylamino)-1-hydroxypropane-1,1-diphosphonic acid, 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid; 1-hydroxy-2-(3-pyridyl)ethane-1,1-diphosphonic acid, 3-[N-(2-phenylthioethyl)-N-methylamino]-1-hydroxypropane-1,1-diphosphonic acid; 1-hydroxy-3-(pyrrolidin-1-yl)propane-1,1-diphosphonic acid, 1-hydroxy-2-(imidazo[1,2-a]pyridin-3-yl)ethane-1,1-diphosphonic acid, a pharmaceutically acceptable salt thereof, and any hydrate thereof.

4. Use according to claim 1 of a methanebisphosphonic acid derivative which is selected from pamidronic acid, alendronic acid, 3-(N-methyl-N-n-pentylamino)-1-hydroxypropane-1,1-diphosphonic acid, 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid, risedronic acid and tiludronic acid, a pharmaceutically acceptable salt thereof, and any hydrate thereof.

5. Use according to claim 1 of a methanebisphosphonic acid derivative which is selected from 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid and 3-amino-1-hydroxypropane-1,1-diphosphonic acid, a pharmaceutically acceptable salt thereof, and any hydrate thereof.

6. Use according to claim 1 of 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid or a pharmaceutically acceptable salt thereof or any hydrate thereof.

7. Use according to claim 1 of 3-amino-1-hydroxypropane-1,1-diphosphonic acid or a pharmaceutically acceptable salt thereof or any hydrate thereof.

8. Use according to claim 1 of disodium pamidronate, or any hydrate thereof.

9. Use according to any one of claims 1 to 8, wherein the prosthesis is a hip prosthesis.

10. Use according to any one of claims 1 to 9, wherein the pharmaceutical composition is adapted to intravenous administration.

11. Use according to any one of claims 1 to 9, wherein the pharmaceutical composition is adapted to transdermal administration.

12. Use according to any one of claims 1 to 9, wherein the pharmaceutical composition is adapted to oral administration.

13. Use of a methanebisphosphonic acid derivative selected from

3-amino-1-hydroxypropane-1,1-diphosphonic acid, 3-(N,N-dimethylamino)-1-hydroxypropane-1,1-diphosphonic acid, 4-amino-1-hydroxybutane-1,1-diphosphonic acid, 6-amino-1-hydroxyhexane-1,1-diphosphonic acid, 3-(N-methyl-N-n-pentylamino)-1-hydroxypropane-1,1-diphosphonic acid, 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid; 1-hydroxy-2-(3-pyridyl)ethane-1,1-diphosphonic acid, and N-methyl pyridinium salts thereof; 1-(4-chlorophenylthio)methane-1,1-diphosphonic acid, 3-[N-(2-phenylthioethyl)-N-methylamino]-1-hydroxypropane-1,1-diphosphonic acid; 1-hydroxy-3-(pyrrolidin-1-yl)propane-1,1-diphosphonic acid, 1-(N-phenylaminothiocarbonyl)methane-1,1-diphosphonic acid, 5-benzoyl-3,4-dihydro-2H-pyrazole-3,3-diphosphonic acid tetraethyl ester, 1-hydroxy-2-(imidazo[1,2-a]pyridin-3-yl)ethane-1,1-diphosphonic acid and 1,1-dichloromethane-1,1-diphosphonic acid; or a pharmaceutically acceptable salt thereof, or any hydrate thereof,

for the prevention and treatment of prosthesis loosening and prosthesis migration in mammals including man.

14. Use according to claim 13 of a methanebisphosphonic acid derivative which is selected from 3-amino-1-hydroxypropane-1,1-diphosphonic acid, 3-(N,N-dimethylamino)-1-hydroxypropane-1,1-diphosphonic acid, 4-amino-1-hydroxybutane-1,1-diphosphonic acid, 6-amino-1-hydroxyhexane-1,1-diphosphonic acid, 3-(N-methyl-N-n-pentylamino)-1-hydroxypropane-1,1-diphosphonic acid, 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid; 1-hydroxy-2-(3-pyridyl)ethane-1,1-diphosphonic acid, and N-methyl pyridinium salts thereof; 1-(4-chlorophenylthio)methane-1,1-diphosphonic acid, 3-[N-(2-phenylthioethyl)-N-methylamino]-1-hydroxypropane-1,1-diphosphonic acid; 1-hydroxy-3-(pyrrolidin-1-yl)propane-1,1-diphosphonic acid, 1-(N-phenylaminothiocarbonyl)methane-1,1-diphosphonic acid, 5-benzoyl-3,4-dihydro-2H-pyrazole-3,3-diphosphonic acid tetraethyl ester, 1-hydroxy-2-(imidazo[1,2-a]pyridin-3-yl)ethane-1,1-diphosphonic acid, a pharmaceutically acceptable salt thereof, and any hydrate thereof.

15. A method of preventing or treating prosthesis loosening and prosthesis migration in mammals, which comprises administering to a mammal in need thereof a therapeutically effective amount of a methanebisphosphonic acid derivative selected from

3-amino-1-hydroxypropane-1,1-diphosphonic acid, 3-(N,N-dimethylamino)-1-hydroxypropane-1,1-diphosphonic acid, 4-amino-1-hydroxybutane-1,1-diphosphonic acid, 6-amino-1-hydroxyhexane-1,1-diphosphonic acid, 3-(N-methyl-N-n-pentylamino)-1-hydroxypropane-1,1-diphosphonic acid, 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid; 1-hydroxy-2-(3-pyridyl)ethane-1,1-diphosphonic acid, and N-methyl pyridinium salts thereof; 1-(4-chlorophenylthio)methane-1,1-diphosphonic acid, 3-[N-(2-phenylthioethyl)-N-methylamino]-1-hydroxypropane-1,1-diphosphonic acid; 1-hydroxy-3-(pyrrolidin-1-yl)propane-1,1-diphosphonic acid, 1-(N-phenylaminothiocarbonyl)methane-1,1-diphosphonic acid, 5-benzoyl-3,4-dihydro-2H-pyrazole-3,3-diphosphonic acid tetraethyl ester, 1-hydroxy-2-(imidazo[1,2-a]pyridin-3-yl)ethane-1,1-diphosphonic acid and 1,1-dichloromethane-1,1-diphosphonic acid; or a pharmaceutically acceptable salt thereof, or any hydrate thereof.

16. A method according to claim 15, wherein the methanebisphosphonic acid derivative administered is selected from 3-amino-1-hydroxypropane-1,1-diphosphonic acid, 3-(N,N-dimethylamino)-1-hydroxypropane-1,1-diphosphonic acid, 4-amino-1-hydroxybutane-1,1-diphosphonic acid, 6-amino-1-hydroxyhexane-1,1-diphosphonic acid, 3-(N-methyl-N-n-pentylamino)-1-hydroxypropane-1,1-diphosphonic acid, 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid; 1-hydroxy-2-(3-pyridyl)ethane-1,1-diphosphonic acid, and N-methyl pyridinium salts thereof; 1-(4-chlorophenylthio)methane-1,1-diphosphonic acid, 3-[N-(2-phenylthioethyl)-N-methylamino]-1-hydroxypropane-1,1-diphosphonic acid; 1-hydroxy-3-(pyrrolidin-1-yl)propane-1,1-diphosphonic acid, 1-(N-phenylaminothiocarbonyl)methane-1,1-diphosphonic acid, 5-benzoyl-3,4-dihydro-2H-pyrazole-3,3-diphosphonic acid tetraethyl ester, 1-hydroxy-2-(imidazo[1,2-a]pyridin-3-yl)ethane-1,1-diphosphonic acid, a pharmaceutically acceptable salt thereof, and any hydrate thereof.

17. A pharmaceutical composition for the prevention and treatment of prosthesis loosening and prosthesis migration which comprises a methanebisphosphonic acid derivative selected from

3-amino-1-hydroxypropane-1,1-diphosphonic acid, 3-(N,N-dimethylamino)-1-hydroxy-

propane-1,1-diphosphonic acid, 4-amino-1-hydroxybutane-1,1-diphosphonic acid, 6-amino-1-hydroxyhexane-1,1-diphosphonic acid, 3-(N-methyl-N-n-pentylamino)-1-hydroxypropane-1,1-diphosphonic acid, 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid; 1-hydroxy-2-(3-pyridyl)ethane-1,1-diphosphonic acid, and N-methyl pyridinium salts thereof; 1-(4-chlorophenylthio)methane-1,1-diphosphonic acid, 3-[N-(2-phenylthioethyl)-N-methylamino]-1-hydroxypropane-1,1-diphosphonic acid; 1-hydroxy-3-(pyrrolidin-1-yl)propane-1,1-diphosphonic acid, 1-(N-phenylaminothiocarbonyl)methane-1,1-diphosphonic acid, 5-benzoyl-3,4-dihydro-2H-pyrazole-3,3-diphosphonic acid tetraethyl ester, 1-hydroxy-2-(imidazo[1,2-a]pyridin-3-yl)ethane-1,1-diphosphonic acid and 1,1-dichloromethane-1,1-diphosphonic acid; or a pharmaceutically acceptable salt thereof, or any hydrate thereof; together with a pharmaceutically acceptable carrier.

18. A pharmaceutical composition according to claim 17, which comprises a methane-bisphosphonic acid derivative selected from 3-amino-1-hydroxypropane-1,1-diphosphonic acid, 3-(N,N-dimethylamino)-1-hydroxypropane-1,1-diphosphonic acid, 4-amino-1-hydroxybutane-1,1-diphosphonic acid, 6-amino-1-hydroxyhexane-1,1-diphosphonic acid, 3-(N-methyl-N-n-pentylamino)-1-hydroxypropane-1,1-diphosphonic acid, 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid; 1-hydroxy-2-(3-pyridyl)ethane-1,1-diphosphonic acid, and N-methyl pyridinium salts thereof; 1-(4-chlorophenylthio)methane-1,1-diphosphonic acid, 3-[N-(2-phenylthioethyl)-N-methylamino]-1-hydroxypropane-1,1-diphosphonic acid; 1-hydroxy-3-(pyrrolidin-1-yl)propane-1,1-diphosphonic acid, 1-(N-phenylaminothiocarbonyl)methane-1,1-diphosphonic acid, 5-benzoyl-3,4-dihydro-2H-pyrazole-3,3-diphosphonic acid tetraethyl ester, 1-hydroxy-2-(imidazo[1,2-a]pyridin-3-yl)ethane-1,1-diphosphonic acid, a pharmaceutically acceptable salt thereof, and any hydrate thereof; together with a pharmaceutically acceptable carrier.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 95/00288

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/66 A61K31/675

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 513 760 (SQUIBB) 19 November 1992 cited in the application ---	
A	US,A,4 711 880 (STAHL ET AL.) 8 December 1987 cited in the application ---	
A	US,A,4 939 130 (JAEGGI ET AL.) 3 July 1990 cited in the application ---	
A	J. BONE MINER. RES., vol. 8, no. suppl.2, 1993 pages S563-S564, P. GOLDBABER 'Strategies to affect bone remodelling.' --- -/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

12 July 1995

Date of mailing of the international search report

21.07.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Klaver, T

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 95/00288

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>CLIN. ORTHOP. RELAT. RES., vol. 287, 1993 pages 61-67, L. RYD ET AL. '99m-Tc-Diphosphonate scintigraphy in successful knee arthroplasty and its relation to micromotion.'</p>	
A	<p>CLIN ORTHOP. RELAT. RES., vol. 167, 1982 pages 106-112, B.J. THOMAS ET AL 'Identification of hip surface arthroplasty failures with TcSC/TcmDP radionuclide imaging.'</p>	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 95/00288

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-513760	19-11-92	JP-A- 5163150	29-06-93
US-A-4711880	08-12-87	AU-B- 591066	30-11-89
		AU-A- 4577785	13-02-86
		CA-A- 1255694	13-06-89
		EP-A, B 0177443	09-04-86
		IE-B- 58077	30-06-93
		JP-C- 1797264	28-10-93
		JP-B- 5008717	02-02-93
		JP-A- 61043196	01-03-86
		KR-B- 9400817	02-02-94
		US-A- 4639338	27-01-87
US-A-4939130	03-07-90	AU-B- 607722	14-03-91
		AU-A- 8145387	26-05-88
		DE-A- 3776880	02-04-92
		EP-A, B 0275821	27-07-88
		HK-A- 114894	27-10-94
		IE-B- 59816	06-04-94
		IL-A- 84497	21-10-94
		JP-A- 63150291	22-06-88
		NO-C- 173446	15-12-93
		ZA-A- 8708698	23-05-88